

# 2

## THE GLOBAL CARBON CYCLE: BIOLOGICAL PROCESSES

Paul G. Falkowski

Department of Earth and Planetary Sciences and Institute of Marine  
and Coastal Sciences, Rutgers University, New Brunswick, NJ 08901, USA

### 2.1 Introduction

Carbon is the fourth most abundant element in our solar system and its chemistry forms the basis of all life on Earth. It is used both as the fundamental building block for all structural biological molecules and as an energy carrier. However, the vast majority of carbon on the surface of this planet is covalently bound to oxygen or its hydrated equivalents, forming mineral carbonates in the lithosphere, soluble ions in the ocean, and gaseous carbon dioxide in the atmosphere. These oxidized (inorganic) forms of carbon are moved on time scales of centuries to millions of years between the lithosphere, ocean and atmosphere via tectonically driven acid-based reactions. Because these reservoirs are so vast (Table 2.1) they dominate the carbon cycle on geological time scales, but because the reactions are so slow, they are also difficult to measure directly within a human lifetime.

The 'geological' or 'slow' carbon cycle is critical for maintaining Earth as a habitable planet (Chapter 2), but entry of these oxidized forms of carbon into living matter requires the addition of hydrogen atoms. By definition, the addition of hydrogen atoms to a molecule is a chemical reduction reaction. Indeed, the addition or removal of hydrogen atoms to and from carbon atoms (i.e., 'redox' reactions), is the core chemistry of life. The processes which drive these core reactions also form a second, concurrently operating global carbon cycle which is biologically catalysed and operates millions of times faster than the geological carbon cycle (Falkowski, 2001). In this chapter, we consider the 'biological', or 'fast' carbon cycle, focusing on how it works, how it evolved, and how it is coupled to the redox chemistry of

a few other elements, especially nitrogen, oxygen, sulfur, and some selected transition metals.

### 2.2 A brief primer on redox reactions

When carbon is directly, covalently linked to hydrogen atoms, the resulting (reduced) molecules are called *organic*. Like acid–base reactions, all reduction reactions must be coupled to a reverse reaction in another molecule or atom; that is the reduction of carbon is coupled to the oxidation of another element or molecule. Under Earth's surface conditions, the addition of hydrogen atoms to carbon requires the addition of energy, while the oxidation of carbon-hydrogen (C–H) bonds yields energy. Indeed the oxidation of C–H bonds forms the basis of energy production for all life on Earth.

Although biologically mediated redox reactions (see Box 2.1) occur rapidly, the products are often kinetically inert. Hence, while it is relatively easy to measure the rate at which a plant converts carbon dioxide into sugars, the product, sugar, is stable. It can be purchased from a local grocery store and kept in a jar in sunlight. It does not spontaneously catch fire or explode. Yet when you eat it, your body extracts the energy from the C–H bonds, and oxidizes the sugar to CO<sub>2</sub> and H<sub>2</sub>O.

### 2.3 Carbon as a substrate for biological reactions

Approximately 75 to 80% of the carbon on Earth is found in an oxidized, inorganic form either as the gas carbon dioxide (CO<sub>2</sub>) or its hydrated or ionic equivalents,

Pools	Quantity (Gt carbon)
Atmosphere	835
Oceans	38,400
Total inorganic	37,400
Surface layer	670
Deep Layer	36,730
Total organic	1,000
Lithosphere	
Sedimentary carbonates	> 60,000,000
Kerogens	15,000,000
Terrestrial biosphere (total)	2,000
Living biomass	600–1,000
Dead biomass	1,200
Aquatic biosphere	1–2
Fossil Fuels	4,130
Coal	3,510
Oil	230
Gas	~300
Other (peat)	250

**Table 2.1** Carbon Pools in the major reservoirs on Earth

namely bicarbonate ( $\text{HCO}_3^-$ ) and carbonate ( $\text{CO}_3^{2-}$ ) (see Fig. 2.1). These inorganic forms of carbon are interconvertible, depending largely on pH and pressure, and the three forms partition into the lithosphere, ocean and atmosphere<sup>1</sup> (see Chapter 3). Virtually all inorganic carbon in the oceans is in the form of  $\text{HCO}_3^-$  with an average concentration of about 2.5 mM. This carbon is removed in association with calcium and magnesium as carbonate minerals. Although the precipitation of carbonates is thermodynamically favourable in the contemporary ocean, it is kinetically hindered, and virtually all carbonates are formed by organisms. The biological precipitation of carbonates is *not* a result of redox reactions, but rather of acid-base reactions; hence, although virtually all carbonates are biologically derived, they remain as oxidized, inorganic carbon. The mineral phases of inorganic carbon are inaccessible to further biological reactions. The total reservoir of inorganic carbon in the ocean is approximately 50 times that of the atmosphere. Indeed, the ocean controls the concentration of  $\text{CO}_2$  in the atmosphere on time scales of decades to millennia.

### 2.3.1 Carbon fixation

Entry of inorganic carbon into biological processes involves a process called carbon ‘fixation’, and there are only two biological mechanisms that lead to the fixation of inorganic carbon: chemoautotrophy and photoautotrophy. Before we consider these in turn, let us first examine what carbon ‘fixation’ is.

<sup>1</sup> Pure forms of carbon as (e.g.) diamond or graphite are relatively rare and do not undergo biological reaction.

The term carbon ‘fixation’ is an anachronism that means ‘to make non-volatile’. It applies when a gaseous  $\text{CO}_2$  is biochemically converted to a solute. There are several enzymatically catalysed reactions that can lead to carbon fixation, however, by far the most important is based on the activity of ribulose 1,5-bisphosphate carboxylase/oxygenase, or Rubisco (Falkowski and Raven, 2007). This enzyme is thought to be the most abundant protein complex on Earth, and it specifically reacts with  $\text{CO}_2$  (i.e., it does not recognize hydrated forms of the substrate). Rubisco catalyses a reaction with a 5 carbon sugar, ribulose 1,5 bisphosphate, leading to the formation of two molecules of 3-phosphoglycerate (see Figs 2.2 and 2.3). This reaction, discovered in the late 1940s and early 1950s by Melvin Calvin, Andrew Benson and Jack Bassham, forms the basis of the pathway of carbon acquisition by most photosynthetic organisms (Benson and Calvin, 1950).

It should be noted that Rubisco imprints a strong biological isotope signature that is used extensively in geochemistry. There are two stable isotopes of C in nature:  $^{12}\text{C}$ , containing 6 protons and 6 neutrons, accounts for 98.90%, and  $^{13}\text{C}$ , containing 6 protons and 7 neutrons, accounts for 1.10%. In the fixation of  $\text{CO}_2$  by Rubisco, the enzyme preferentially reacts with the lighter isotope; the net result is that 3-phosphoglycerate is enriched by about 25 parts per thousand in  $^{12}\text{C}$  relative to the  $\text{CO}_2$  in the air or water (Park, 1961). This isotopic fractionation provides a basis for understanding the impact of the biological carbon cycle over geological time (Kump and Arthur, 1999).


The fixation of  $\text{CO}_2$  by Rubisco is *not* an oxidation/reduction reaction; the carboxylic acid group has the same oxidation state as  $\text{CO}_2$ . The biochemical

**Box 2.1** Redox reactions

The term *oxidation* was originally used by chemists in the latter part of the 18th century to describe reactions involving the addition of oxygen to metals, forming metallic oxides. For example:



The term *reduction* was used to describe the reverse reaction, namely the removal of oxygen from a metallic oxide, for example, by heating with carbon:



Analysis of these reactions established that the addition of oxygen is accompanied by the removal of electrons from an atom or molecule. Conversely, reduction is accompanied by the addition of electrons. In the specific case of organic reactions that involve the reduction of carbon, the addition of electrons is usually balanced by the addition of protons. For example, the reduction of carbon dioxide to formaldehyde requires the addition of four electrons *and* four  $\text{H}^+$  – that is, the equivalent of four hydrogen atoms.



Thus, from the perspective of organic chemistry, oxidation may be defined as the addition of oxygen, the loss of electrons, or the loss of hydrogen atoms (but not hydrogen ions,  $\text{H}^+$ ); conversely, reduction can be defined as the removal of oxygen, the addition of electrons, or the addition of hydrogen atoms.

Oxidation–reduction reactions only occur when there are pairs of substrates, forming pairs of products:



Photosynthesis uses energy from the sun to reduce inorganic carbon to form organic matter; i.e., photosynthesis is a biochemical reduction reaction. In oxygenic photosynthesis,  $\text{CO}_2$  is the recipient of the electrons and protons, and thus becomes reduced (it is the A in Equation B2.1.4). Water is the electron and proton donor, and thus becomes oxidized (it is the B in Equation B2.1.4). The oxidation of two moles of water requires the addition of 495 kJ of energy. The reduction of  $\text{CO}_2$  to the simplest organic carbon molecule, formaldehyde, requires 176 kJ of energy. The energetic efficiency of photosynthesis can be calculated by dividing the energy stored in organic matter by that required to split water into molecular hydrogen and oxygen. Thus, the maximum overall efficiency of photosynthesis, assuming no losses at any intermediate step, is 176/495 or about 36%.

*reduction* of 3-phosphoglycerate is the second step in the carbon fixation pathway, and leads to the formation of an aldehyde. This is the only reduction step in the so-called Calvin cycle. The rest of the pathway is primarily devoted to regenerating ribulose 1,5-bisphosphate, and leaves no discernable geochemical signal.


## 2.3.2 Chemoautotrophy

Chemoautotrophs (literally, ‘chemical self feeders’) are organisms capable of reducing sufficient inorganic carbon to grow and reproduce in the absence of light energy and without an external organic carbon source. Chemoautotrophs likely evolved very early in Earth’s history and this process is exclusively carried out by prokaryotic organisms in both the domains Archaea and Bacteria (Stevens and McKinley, 1995).

Early in Earth’s history,  $\text{H}_2$  was probably an important constituent of the atmosphere a major reductant used by organisms to reduce inorganic carbon to organic biomass (Jørgensen, 2001). Although this process can still be found in marine sediments, where  $\text{H}_2$  is produced during the anaerobic fermentation of organic matter, and in hydrothermal environments, where the gas is produced as a byproduct of serpentinization, free  $\text{H}_2$  is scarce on Earth’s surface. Rather, most of the hydrogen is combined (oxidized) by microbes with other atoms, such as sulfur or oxygen, and to a much lesser extent, nitrogen. Hence, most contemporary chemoautotrophs oxidize  $\text{H}_2\text{S}$  or  $\text{NH}_4^+$ , but also other reduced compounds such as  $\text{Fe}^{2+}$ .

The driver for chemoautotrophic carbon fixation ultimately depends on a thermodynamically favourable redox gradient. For example, the oxidation of  $\text{H}_2\text{S}$  by microbes in deep sea vents is coupled to the reduction of oxygen in the surrounding water. Hence, this reaction is dependent on the chemical redox gradient between the ventilating mantle plume and the ocean interior that thermodynamically favours oxidation of the plume gases (Jannasch and Taylor, 1984) (Box 2.2).

Chemoautotrophy supplies a relatively small amount of organic carbon to the planet (probably <1%), however this mode of nutrition is critically important in sediments, anoxic basins, and in completing several elemental cycles, including that of N and S. Thus, chemoautotrophy is common in sediments and anoxic water columns where strong redox gradients develop. A classic example would be the oxygen-sulfide interface in microbial mats where sulfide-oxidizing chemoautotrophs thrive, although countless other examples could also be named (Canfield and Raiswell, 1999). Reductants for chemoautotrophs can be generated within in the Earth’s crust. An important example, as mentioned above, is the hydrothermal fluids generated in mid-ocean ridge spreading centres. Here, the sulfide and ferrous iron liberated with the fluids sup-



**Figure 2.1** The relative distribution of the three major species of dissolved inorganic carbon in water as a function of pH. Note that at pH of seawater (~8.1), approximately 95% of the inorganic carbon is in the form of bicarbonate anion.


port the chemoautotrophic growth of sulfide- and Fe-oxidizers, which use oxygen as the oxidant.

However, early in Earth's history, due to the lack of oxygen, the redox gradients would have been small and hence there would have been no pandemic outbreak of chemoautotrophy. The vents themselves would have supplied H<sub>2</sub> and CO<sub>2</sub>, for example, which could have supported the chemoautotrophic growth of methanogens (Canfield *et al.*, 2006), but this would have been on a much smaller scale than the chemoautotrophic sulfide oxidation supported in modern vents. Importantly, magma chambers, volcanism, and vent fluids are tied to either subduction or to spreading regions, which are transient features of Earth's crust and hence only temporary habitats for chemoautotrophs. In the Archean and early Proterozoic oceans, the chemoautotrophs would have had to have been dispersed throughout the oceans by physical mixing in order to colonize new vent regions (Raven and Falkowski, 1999).


### 2.3.3 Photoautotrophy

Photoautotrophy ('self feeding on light') is the biological conversion of light energy to the fixation of CO<sub>2</sub> in the form of organic carbon compounds. To balance the electrons, a source of reductant is also required. It should be noted that while all photoautotrophs are photosynthetic, not all photosynthetic organisms are photoautotrophs. Many organisms are capable of photosynthesis but can (and, sometimes must) supplement that metabolic strategy with the assimilation of organic carbon (Falkowski and Raven, 2007).


Photoautotrophy can be written as an oxidation-reduction reaction of the general form:



In this representation, light is specified as a substrate, with some of the energy of the absorbed light stored in the products. All photoautotrophic bacteria, with the important exception of the cyanobacteria, are incapable of evolving oxygen. In these (mostly) anaerobic organisms, compound A is, for example, an atom of sulfur, and the pigments are bacteriochlorophylls (Van Niel, 1941; Blankenship *et al.*, 1995). All other photoautotrophs, including the cyanobacteria, eukaryotic algae, and higher plants, are oxygenic; that is, Equation 2.1 can be modified to:




where Chl *a* is the ubiquitous plant pigment chlorophyll *a*. Equation 2.2 implies that chlorophyll *a* catalyses a reaction or a series of reactions whereby light energy is used to oxidize water:




yielding gaseous, molecular oxygen. True photoautotrophy is restricted to the domains Bacteria and Eukarya. Although some Archaea and Bacteria use the pigment rhodopsin to harvest light, they require organic matter to fuel their metabolism (Beja *et al.*, 2000) and are not photoautotrophs.

## 2.4 The evolution of photosynthesis

We have discussed above the production of organic matter by photoautotrophy, but photosynthesis is more




Structures of key intermediates



**Figure 2.2** The reactions of ribulose 1,5 bisphosphate carboxylase/oxygenase. In the carboxylation process, the enzyme reacts with CO<sub>2</sub> to produce 2 molecules of 3-phosphoglycerate, which is the first stable intermediate in the Calvin-Benson cycle. Alternatively, in the oxygenase process, the enzyme reacts with O<sub>2</sub>, leading to the production 2 phosphoglycolate which ultimately is respired to CO<sub>2</sub>.

broadly defined than this. Normally we consider photosynthesis to include all organisms that use light energy to synthesize new cells and this includes also photoheterotrophs which incorporate organic matter from the environment into their biomass. By far the most efficient and ubiquitous light harvesting systems for photosynthesis are based on porphyrins. The metabolic pathway for the synthesis of porphyrins is extremely old (Mauzerall,

1992); relic porphyrin molecules can be isolated from some ancient Archean (older than 2.5 billion years) rocks. It has been proposed that the porphyrin-based photosynthetic energy conversion apparatus originally arose from the need to prevent UV radiation from damaging essential macromolecules such as nucleic acids and proteins (Mulikidjanian and Junge, 1997). Indeed, most photosynthetic bacteria retain an ability to harvest UV light for



**Figure 2.3** The basic Calvin–Benson cycle. CO<sub>2</sub> is condensed with ribulose 1,5 bisphosphate to form 3 phosphoglycerate (1), which is subsequently reduced to an aldehyde (reactions 2 and 3). The remaining reactions in the cycle are directed towards regenerating ribulose 1,5 bisphosphate; the intermediate structures are indicated.

### Box 2.2 The Nernst Equation

As mentioned in the text, oxidation–reduction reactions involve the transfer of electrons. The tendency for a molecule to accept or release an electron is viewed relative to the ability of a ‘standard’ molecule to do the same, and this is normally the standard hydrogen electrode (SHE), which is represented by the following reaction:



The SHE is defined at 25 °C, and one atmosphere of H<sub>2</sub> gas, and for an H<sup>+</sup><sub>(aq)</sub> activity of 1 (pH = 0). The tendency of a species to accept or liberate electrons is formally known as *electrode potential* or *redox potential*, *E*. The SHE is arbitrarily assigned an *E* of 0. The redox potential of reactions at standard state (unit activity for reactants and products), and relative to the SHE are given the designation *E*<sub>0</sub>. In comparing to the SHE, reactions are always written as reduction reactions after the following general form:



It is often more useful to define the redox potential at pH = 7, an environmentally relevant pH, and when so defined, the redox potential is denoted by the symbols *E*<sub>0</sub>' or sometimes *E*<sub>m7</sub>'. The *E*<sub>0</sub>' for a standard hydrogen electrode is –420 mV.

It is rare for organisms to live under standard-state chemical conditions (unit activity concentrations or reactants and products), and the electrode potential can be modified with the Nernst equation to reflect environmental conditions. With the general equation represented in B2.2.2, the Nernst equation is written as:

$$E = E_0 + 2.3(RT/nF) \log_{10} (a^a_{\text{Aoxid}} a^c_{\text{H}^+} / a^d_{\text{Ared}} a^g_{\text{G}}) \quad (\text{B2.2.3})$$

Where *E* is the redox potential (in volts) under environmental conditions, *E*<sub>0</sub> is the standard redox potential, *F* is Faraday's constant (= 96 485 coulombs = the electrical charge in 1 mole of electrons), *n* is the number of moles of electrons (Faradays) transferred in the half-cell reaction, *R* is the Boltzmann gas constant, *T* is temperature in Kelvin, and *a*<sub>*y*</sub><sup>*x*</sup> represents the activity (sometimes concentration is used, but this is not strictly correct) of species *y* raised to the stoichiometric factor *x* in the balanced half reaction equation. The value of 2.3(*RT*/*F*) is 59 mV.

photosynthesis, but these bacteria cannot split water and evolve oxygen. These are termed 'anoxygenic' organisms, with light-mediated sulfide oxidation, as introduced above, a common (but not exclusive) metabolism.

In the ancient oceans, anoxygenic photosynthesis had profound biogeochemical consequences. It led not only to the formation of organic matter, but to the oxidation of such reductants as  $\text{Fe}^{2+}$  (to  $\text{Fe}^{3+}$ , which precipitated as Fe oxides), which was found in abundance in ancient oceans and  $\text{S}^{2-}$  (to  $\text{S}^0$  or  $\text{SO}_4^{2-}$ ), which was found in ancient hydrothermal springs, in ancient oceans during some time periods (Canfield, 1998; Brocks *et al.*, 2005) and in contemporary analogs such as in Yellowstone National Park. While these reductants were ultimately resupplied via weathering in the case of Fe, or from hydrothermal fluids in the case of sulfide, the rate of supply was slow relative to the rate of at which organisms can photosynthesize. Hence there was a drive to find a reductant for photosynthesis that is virtually limitless. The obvious molecule is  $\text{H}_2\text{O}$ .

Liquid water contains  $\sim 55 \text{ kmol H}_2\text{O per m}^3$ , and there are  $10^{18} \text{ m}^3$  of water in the hydrosphere and cryosphere. However, the use of  $\text{H}_2\text{O}$  as a reductant for  $\text{CO}_2$ -fixation to organic matter requires a larger energy input than does the use of  $\text{Fe}^{2+}$  or  $\text{S}^{2-}$ . Indeed, for oxygenic photosynthesis to occur, several innovations on the old anoxygenic anaerobic photosynthetic machinery had to occur (Blankenship *et al.*, 2007). Among these innovations were the evolution of: (a) a new photosynthetic pigment, chlorophyll *a*, which operates at a higher energy level than bacterial chlorophylls; (b) two photochemical reaction centres that operate in series, one of which splits water, the second of which forms a biochemical reductant that is used to reduce inorganic carbon; and (c) a unique complex comprised of four Mn atoms bound to a group of proteins that forms the 'oxygen evolving complex', i.e., the site in which four electrons are sequentially extracted from two molecules of liquid water, one at a time, via the absorption of four photons.

The photochemical apparatus responsible for oxygenic photosynthesis is the most complex energy transduction system found in nature; there are well over 100 genes necessary for its synthesis (Shi and Falkowski, 2008). It appears to have arisen only once, in a single clade of bacteria (the cyanobacteria), and has never been appropriated by any other prokaryote. The origin and evolutionary trajectory of oxygenic photosynthesis remains obscure (Falkowski and Raven, 2007). It almost certainly arose sometime in the Archean Eon, although the timing is uncertain (see Chapter 7). The two photosystems appear to have different origins: the water splitting system is derived from purple photosynthetic bacteria, while the second reaction centre is derived from green sulfur bacteria. How the two reactions became incorporated into a single organism is unknown.

In order to oxidize water, the photochemical reaction must generate an oxidant with a potential of  $+0.8 \text{ V}$  ( $E_m$ ) or more. This is significantly greater than is found in any extant anoxygenic photoautotroph (the highest is ca.  $+0.4 \text{ V}$ ). The oxidizing potential in oxygenic photosynthesis is the highest in nature, and ultimately, oxygenic photosynthesis became the primary mechanism for reducing  $\text{CO}_2$  and forming organic carbon. Once established, it freed the microbial world from a limited supply of reductants for carbon fixation, and it decoupled the biological carbon cycle from the geological carbon cycle on time scales of millenia (Falkowski and Raven, 2007).

## 2.5 The evolution of oxygenic phototrophs

### 2.5.1 *The cyanobacteria*

Cyanobacteria are the only oxygenic phototrophs known to have existed before  $\sim 2 \text{ Ga}$ . There is some suggestion that the 1.8 billion year old fossil *Grypania* may represent an eukaryotic algae (Han and Runnegar, 1992), but this has not been firmly established. Cyanobacteria numerically dominate the phototrophic community in contemporary marine ecosystems, and clearly their continued success bespeaks an extraordinary adaptive capacity.

By 2 billion years ago, cyanobacteria were probably the major primary producers (a primary producer is an organism that supplies organic matter to heterotrophs), with likely contributions from anoxygenic phototrophs and chemoautotrophs. In the contemporary ocean, the cyanobacteria fix approximately 60% of the  $\sim 45 \text{ Pg C}$  assimilated annually by aquatic phototrophs (Falkowski and Raven, 2007). Their proportional contribution to 'local' marine primary productivity is greatest in the oligotrophic central ocean gyres that form about 70% of the surface waters of the seas. Two major groups of marine cyanobacteria can be distinguished. The phycobilin-containing *Synechococcus* are more abundant nearer the surface and the (divinyl) chlorophyll *b*-containing *Prochlorococcus* are generally more abundant at depth (Chisholm, 1992) (Table 2.2).

Some cyanobacteria not only fix inorganic carbon, but also fix  $\text{N}_2$ . Biological reduction of  $\text{N}_2$  to  $\text{NH}_3$  (i.e., 'fixation') is catalysed by nitrogenase, a heterodimeric enzyme that is irreversibly inhibited by  $\text{O}_2$ . Molecular phylogenetic trees suggest that  $\text{N}_2$  fixation evolved in Bacteria prior to the evolution of oxygenic photosynthesis (Zehr *et al.*, 1997) and was acquired by cyanobacteria relatively late in their evolutionary history (Shi and Falkowski, 2008). The early evolution of nitrogenase is also indicated by the very large Fe requirement for the enzyme; the holoenzyme contains 38 iron atoms. This transition metal was much more available in the water column of the oceans in the Archean and

	Ocean NPP		Land NPP
Seasonal			
April–June	10.9		15.7
July–September	13.0		18.0
October–December	12.3		11.5
January–March	11.3		11.2
Biogeographic			
Oligotrophic	11.0	Tropical rainforests	17.8
Mesotrophic	27.4	Broadleaf deciduous forests	1.5
Eutrophic	9.1	Broadleaf and needleleaf forests	3.1
Macrophytes	1.0	Needleleaf evergreen forests	3.1
		Needleleaf deciduous forest	1.4
		Savannas	16.8
		Perennial grasslands	2.4
		Broadleaf shrubs with bare soil	1.0
		Tundra	0.8
		Desert	0.5
		Cultivation	8.0
Total	48.5		56.4

**Table 2.2** Annual and seasonal net primary production (NPP) of the major units of the biosphere (after Field *et al.*, 1998)

Source: Field *et al.* (1998). All values in GtC. Ocean NPP estimates are binned into three biogeographic categories on the basis of annual average  $C_{\text{sat}}$  for each satellite pixel, such that oligotrophic =  $C_{\text{sat}} < 0.1 \text{ mg m}^{-3}$ , mesotrophic =  $0.1 < C_{\text{sat}} < 1 \text{ mg m}^{-3}$ , and eutrophic =  $C_{\text{sat}} > 1 \text{ mg m}^{-3}$  (Antoine *et al.*, 1996). This estimate includes a 1 GtC contribution from macroalgae (Smith, 1981). Differences in ocean NPP estimates between Behrenfeld and Falkowski (1997b) and those in the global annual NPP for the biosphere and this table result from (i) addition of Arctic and Antarctic monthly ice masks; (ii) correction of a rounding error in previous calculations of pixel area; and (iii) changes in the designation of the seasons to correspond with Falkowski *et al.* (1998). The macrophyte contribution to ocean production from the aforementioned is not included in the seasonal totals. The vegetation classes are those defined by DeFries and Townshend (DeFries and Townshend, 1994).

early Proterozoic Eons than it is today (Berman-Frank *et al.*, 2001). Indeed, iron has been suggested to limit nitrogen fixation in the contemporary ocean (Falkowski 1997).

### 2.5.2 The eukaryotes

Eukaryotes are unicellular or multicellular organisms whose cells possess nuclei that contain their DNA. All eukaryotes were derived from a symbiotic association between a host cell closely affiliated with the Archaea (Woese, 1998), and an organelle derived from the Bacteria. The symbiosis occurred via the engulfment of the latter cell by the former, after which most of the genes in the bacterium were either lost or transferred to the host. There appears to have been only two primary symbiotic events. The first involved a purple bacterium, which may have been photosynthetic (Woese *et al.*, 1984); this organism ultimately came to be a mitochondrion, the primary energy producing system for eukaryotes. In a second symbiotic event, a host eukaryote engulfed a cyanobacterium. The cyanobacterium would, over time, lose many of its genes and become a plastid (or,

specifically in green plants, a chloroplast) (Martin and Herrmann, 1998), leading the evolution of an oxygenic photosynthetic eukaryote. Because they lost so many genes, the endosymbiotic cyanobacteria became incapable of independent existence outside the symbiotic association; essentially they are enslaved by the host cell (Bhattacharya and Medlin, 1998). All eukaryotic photoautotrophs are oxygenic (Falkowski *et al.*, 2004).

While there are eight major phyla of photosynthetic eukaryotes, they can be broadly lumped into two major groups: a green and a red clade. The former contain, in addition to chlorophyll *a*, a second photosynthetic pigment, chlorophyll *b*. The latter does not contain chlorophyll *b*, but rather other pigments, especially chlorophyll *c*. Regardless of the evolutionary history of the host, however, the core photosynthetic machinery is highly conserved. That is, the same basic structure that evolved in cyanobacteria is found in every oxygenic photosynthetic eukaryote. Indeed, in some cases, the gene sequences are so highly conserved across the ca. 3 billion years of evolution that they have been called ‘frozen metabolic accidents’ – genes that essentially have ceased evolving (Shi *et al.*, 2005).



Approximately 500 million years ago, one green alga ultimately became the progenitor of all higher plants (Knoll, 1992; Kenrick and Crane, 1997). Although there are in excess of 250,000 species of extant higher plants, they are all relatively closely related to each other. Except for diatoms that live in soils, member of the red clade of photosynthetic eukaryotes successfully colonized land. These organisms, however, became extremely successful in aquatic ecosystems. All of the major photosynthetic eukaryotes in the ocean arose by *secondary* endosymbiosis, where eukaryotic phagotrophic flagellates ingested eukaryotic phototrophs (Falkowski *et al.*, 2004). This process led to the evolution of diatoms and other closely related groups (including the kelps), haptophyte algae including coccolithophores, and dinoflagellates (Delwiche, 2000). These three groups, which are very distantly related, have left fossils that allow some physical reconstruction of their evolutionary history. There is fossil evidence that a (macrophyte) red alga existed 1.2Ga, with unicellular (cysts) and macrophytic green algae from 1 and 0.6Ga bp respectively.

While fossils referable to heterokonts (which include diatoms) are known from the late Proterozoic, it is possible that these are not photosynthetic, granted the much later (280Ma bp) origin of *photosynthetic* heterokonts suggested by molecular clock data calibrated from the molecular phylogeny and fossil record of diatoms (Bhattacharya *et al.*, 1992; Kooistra *et al.*, 2007). The precursors of modern dinoflagellates probably originated more than 250 million years ago (Falkowski *et al.*, 2004), while at least the coccolithophorid variants of the haptophytes are first known from fossils of the early Triassic period (de Vargias *et al.*, 2007).

This brief account of the evolution and diversification of aquatic photoautotrophs reveals that there is much more higher-taxon diversity among aquatic than terrestrial primary producers; 95% or more of species involved in terrestrial primary producers are embryophytes derived from a single class (Charophyceae) of a single Division (Chlorophyta) among the eight Divisions of eukaryotic aquatic photoautotrophs. By contrast, there is no comparable taxonomic dominance of primary producers in aquatic ecosystems. Simply put, the land is green and the oceans are red (Falkowski *et al.*, 2003).

## 2.6 Net primary production


Photosynthesis allows an organism to convert inorganic carbon to organic matter, yet the organism must also use some of that organic matter for its own metabolic demands. Hence, ecologists have devised another term: net primary production (NPP), defined as that fraction of photosynthetically produced carbon that is retained following all respiratory costs of the photoautotrophs

and consequently is available to the next trophic level in the ecosystem (Lindeman, 1942).

In terrestrial ecosystems, NPP is derived from empirical measurements of leaf area (i.e., the ability of a plant to absorb light), leaf nitrogen content (a surrogate for the content of the photosynthetic machinery, as leaves are the photosynthetic organ of a plant), irradiance, temperature and water availability. To a first order, the distribution of plants on land is controlled by water availability (Mooney *et al.*, 1987). When water is available, the productivity of terrestrial plants is mainly controlled by how much light is intercepted by the leaves (Nobel *et al.*, 1993; Field *et al.*, 1998). Models for NPP on land are relatively easy to verify; plants can be weighed and their carbon content can be measured.

In aquatic ecosystems, net primary production is generally estimated from incorporation rates of  $^{14}\text{C}$ -labelled inorganic carbon (supplied as bicarbonate) into organic matter. Alternative approaches include oxygen exchange in an enclosed volume (a technique which is far less sensitive and much more time-consuming) and variable fluorescence (which embraces a variety of approaches) (Falkowski *et al.*, 1998). As for terrestrial ecosystems, extrapolation of photosynthetic rates from any technique to the water column requires a mathematical model. The commonly used strategy is to derive an empirical relationship between photosynthesis and irradiance for a given phytoplankton or macrophyte community, and to integrate that (instantaneous) relationship over time and depth in the water column. Assuming some respiratory costs during the photo-period and during the dark, this approach gives an estimate of net primary production for the water column (Behrenfeld and Falkowski, 1997).

Using satellite-based images of terrestrial vegetation and ocean water leaving radiances (i.e., the visible light scattered back to space from the ocean), it is possible to construct maps of the global distributions of NPP. Briefly, the terrestrial NPP is constructed from estimates of leaf cover and irradiance. For the oceans, phytoplankton chlorophyll is derived based on the ratio of green to blue light that is reflected back to space. In the absence of photosynthetic pigments in the upper ocean, radiances leaving water would appear blue to an observer in space. In the presence of chlorophylls and other photosynthetic pigments, some of the blue radiation is absorbed in the upwelling stream of photons, thereby reducing the overall radiance; the water becomes 'darker'. The fraction of blue light that is absorbed is proportional to the chlorophyll concentration. After accounting for scattering and absorption by the atmosphere, satellite-based maps of both terrestrial vegetation and ocean color can be used to infer the spatial and temporal variations of photosynthetic biomass from which NPP is derived (Behrenfeld and Falkowski, 1997).



**Figure 2.4** Seasonal, maps of global inventories of photosynthetic biomass in the oceans and on land as derived from satellite images. These data are used to construct global estimates of primary production (e.g. Field *et al.*, 1998). Note the strong seasonal variations in biomass, especially at high latitudes in both hemispheres (data courtesy of NASA).

Monthly, seasonal and annual maps of global ocean net primary production have been developed that incorporate a primary production model into global ocean colour images of phytoplankton chlorophyll (Fig. 2.4). Such models suggest that annual net ocean photosynthetic carbon fixation is about  $45 \pm 5 \text{ GtC}$  per annum (Behrenfeld and Falkowski, 1997a, b). This productivity is driven by a photosynthetic biomass that amounts to ca.  $1 \text{ GtC}$ . Hence, phytoplankton biomass in the oceans

turns over on the order of once every two weeks; the overwhelming majority ( $\sim 85\%$ ) of the productivity is consumed by heterotrophs in the upper ocean, and about 15% sinks into the ocean interior where virtually all is consumed ('remineralized') and converted back to inorganic carbon (see also Chapter 3). Only a very small fraction of the organic matter produced is buried in marine sediments. Simply put, less than 1% of the photosynthetic biomass on Earth accounts for about 50% of

the net primary productivity (Field *et al.*, 1998). What about the other 50%?

Net primary production in terrestrial ecosystems accounts for approximately 52 GtC per annum, and is supported by a biomass of approximately 500 GtC (Field *et al.*, 1998). Thus, the turnover time of terrestrial plant biomass is on the order of a decade. As in the ocean, the overwhelming majority of terrestrial plant production is recycled back to inorganic carbon, primarily by fungi and microbes which degrade dead plants. In terrestrial ecosystems, virtually all of the photosynthetic biomass is consumed by heterotrophs long after it was produced. For example, the leaves on deciduous trees remain until the autumn, when they then die and fall to the ground. Over the coming years, virtually 100% of the leaf litter will be consumed by microbes and fungi, but very little will have been eaten by an animal. Terrestrial ecosystems contain large amounts of carbon associated with living structures (e.g. wood in trees), but their photosynthetic rates are not sufficiently different than aquatic ecosystems, which have a much faster rate of ecological metabolism. Thus, although the recycling time for terrestrial plant production is much longer than that in the ocean, the fates are basically the same; virtually all of the organic matter is reprocessed by heterotrophs resulting in a biological cycle that has relatively small impact on the ocean/atmosphere inventory of CO<sub>2</sub> on time scales of decades or centuries (Falkowski *et al.*, 2000).

## 2.7 What limits NPP on land and in the ocean?

All organisms are primarily composed of six major elements: H, C, N, O, P and S (Schlesinger, 1997). The ratios of these elements can vary widely, depending on the type of organism and the ecosystem; however, one or more of these elements often limits NPP. The limitations fundamentally differ between aquatic and terrestrial ecosystems.

Terrestrial plants have an absolute requirement for water, which clearly is not a problem for aquatic photoautotrophs. However, even when water is ample, carbon fixation is not necessarily maximal. One of the major limiting factors limiting NPP on land is the availability of CO<sub>2</sub> itself. Although Rubisco is the most abundant protein on Earth, it has a low affinity for CO<sub>2</sub>. Saturation values for CO<sub>2</sub> in terrestrial ecosystems are in the order of ca. 500 parts per million, significantly above that found in the contemporary atmosphere (Mooney *et al.*, 1991). To make matters even worse, the enzyme also 'mistakes' O<sub>2</sub> as a substrate (hence, the origin of the term 'oxygenase' in the appellation of the enzyme). Indeed, in most terrestrial plants, approximately 35% of the time, the enzyme reacts with O<sub>2</sub> leading to the imme-

diately respired of two carbon products which are never converted into biomass. This 'photorespiratory' process leads to a large loss in potential NPP when CO<sub>2</sub> is not saturating. Some terrestrial plants have overcome this limitation by developing a high-affinity carbon fixation process which allows them to use lower levels of CO<sub>2</sub>; these so-called C<sub>4</sub> plants are primarily tropical grasses (such as cane sugar), and account for approximately 15% of global terrestrial NPP (Berry, 1999). Thus, on land, both water and CO<sub>2</sub> are major factors limiting NPP.

In aquatic ecosystems, water and inorganic carbon are not limiting. Although the concentration of gaseous CO<sub>2</sub> in the oceans is only ~10 μM, the total concentration of inorganic carbon is ~2.5 mM, of which 95% is in the form of HCO<sub>3</sub><sup>-</sup>. Although HCO<sub>3</sub><sup>-</sup> cannot be used by Rubisco directly, the anion can be transported into cells from the sea and converted to CO<sub>2</sub> by the enzyme carbonic anhydrase, which is one of the most catalytically active enzymes known. The transport and dehydration of HCO<sub>3</sub><sup>-</sup> by aquatic photoautotrophs is called a 'carbon concentrating mechanism', and it virtually assures that carbon never limits NPP in aquatic ecosystems (Kaplan and Reinhold, 1999). If water or inorganic carbon are not limiting, what is?

Globally, nitrogen and phosphorus are the two elements that immediately limit NPP in both lakes and the oceans. It is frequently argued that since N<sub>2</sub> is abundant in both the ocean and the atmosphere, and since in principle it can be biologically reduced to the equivalent of NH<sub>3</sub> by N<sub>2</sub>-fixing cyanobacteria, then nitrogen cannot be limiting on geological time scales (Redfield, 1958; Barber, 1992). Therefore, phosphorus, which is supplied to the ocean by the weathering of continental rocks, must ultimately limit biological productivity (Broecker and Peng, 1982; Tyrell, 1999). The underlying assumptions of these tenets should, however, be considered within the context of the evolution of biogeochemical cycles.

The main source of fixed inorganic nitrogen to the oceans is biological nitrogen fixation. This is strictly a prokaryotic process, and in the modern ocean, it is conducted mainly by cyanobacteria. On the early Earth, however, before the evolution of nitrogen fixers, electrical discharge or bolide impacts may have promoted NO formation from the reaction between N<sub>2</sub> and CO<sub>2</sub>, but the yield for these reactions is extremely low. Moreover, any volcanogenic NH<sub>3</sub> in the atmosphere would have photodissociated from UV radiation (Kasting, 1990) while N<sub>2</sub> would have been stable (Warneck, 1988; Kasting, 1990). Biological N<sub>2</sub> fixation is a strictly anaerobic process (Postgate, 1971), and the genes encoding the catalytic subunits for nitrogenase are highly conserved in cyanobacteria and other prokaryotes that fix nitrogen, strongly suggesting a common ancestral origin (Zehr *et al.*, 1995). Since fixed inorganic nitrogen was likely to

have been scarce prior to the evolution of diazotrophic (nitrogen fixing) organisms, there was strong evolutionary selection for nitrogen fixation on the early Earth.

The formation of nitrate from ammonium by nitrifying bacteria requires molecular oxygen (see Chapter 7); hence, nitrification must have evolved following the formation of free molecular oxygen in the oceans by oxygenic photoautotrophs. Therefore, from a geological perspective, the conversion of ammonium to nitrate probably proceeded rapidly and provided a substrate,  $\text{NO}_3^-$ , that eventually could serve both as a source of nitrogen for photoautotrophs and as an electron acceptor for a diverse group of heterotrophic, anaerobic bacteria, the denitrifiers.

In the sequence of the three major biological processes that constitute the nitrogen cycle, denitrification must have been the last to emerge. This process, which permits the reduction of  $\text{NO}_3^-$  to (ultimately)  $\text{N}_2$ , requires hypoxic or anoxic environments and is sustained by high sinking fluxes of organic matter. With the emergence of denitrification, the ratio of fixed inorganic N to dissolved inorganic phosphate in the ocean interior could only be depleted in N relative to the sinking flux of the two elements in POM. Indeed, in all of the major basins in the contemporary ocean, the N:P ratio of the dissolved inorganic nutrients in the ocean interior is conservatively estimated at 14.7 by atoms (Fanning, 1992) or less (Anderson and Sarmiento, 1994), compared to the Redfield ratio of 16/1.

There are three major conclusions that may be drawn from the foregoing discussion. First, because the sinking flux of particulate matter has an organic N to P ratio (about 16/1) which exceeds the N to P ratio of the deep ocean dissolved inorganic pool (average 14.7/1 as noted above), the average upward flux of nutrients is slightly enriched in P relative to the requirements of the photoautotrophs (Redfield, 1958; Gruber and Sarmiento, 1997). Hence, with some exceptions (Wu *et al.* 2000), dissolved inorganic fixed nitrogen generally limits primary production throughout most of the world's oceans on ecological times scales (Barber, 1992); (Falkowski, 1998). Second, the N to P ratio of the deep-ocean inorganic dissolved pool was established by biological processes, not vice versa (Redfield, 1934; Redfield *et al.*, 1963). Third, if dissolved inorganic nitrogen rather than phosphate limits productivity in the oceans, then it follows that the ratio of nitrogen fixation to denitrification plays a critical role in determining primary production and the net biologically mediated exchange of  $\text{CO}_2$  between the atmosphere and ocean (Codispoti, 1995).

On ecological time scales, primary production is limited by nutrient supply and the efficiency of nutrient utilization in the euphotic zone. There are three major areas of the world ocean where inorganic nitrogen and phosphate are in excess throughout the year, yet the

mixed layer depth appears to be shallow enough to support active primary production (if the mixed layer depth extends below the depth where sufficient light is available for primary production, production becomes limited): these are the eastern equatorial Pacific, the subarctic Pacific, and Southern (i.e., Antarctic) Oceans. For the subarctic North Pacific, Miller *et al.* (1991) suggested a tight coupling between phytoplankton production and consumption by zooplankton. This grazer-limited hypothesis was used to explain why the phytoplankton in the North Pacific do not form massive blooms in the spring and summer like their counterparts in the North Atlantic (Banse, 1992).

In the mid-1980s, however, it became increasingly clear that the concentration of trace metals, especially iron, was extremely low in all three of these regions (Martin *et al.*, 1991). Indeed, in the eastern equatorial Pacific the concentration of soluble iron in the euphotic zone is only 100 to 200 pM, about an order of magnitude lower than found in other areas of the open ocean. Although iron is the most abundant transition metal in the Earth's crust, in its most commonly occurring form,  $\text{Fe}^{3+}$ , it is virtually insoluble in oxygenated water. The major source of iron to the euphotic zone of the ocean is aeolian dust, originating from continental deserts. In the three major areas of the world oceans with high inorganic nitrogen in the surface waters and low chlorophyll concentrations, the flux of aeolian iron is extremely low (Duce and Tindale, 1991). In experiments in which iron was artificially added on a relatively large scale to the waters in the equatorial Pacific, Southern Ocean, and subarctic Pacific there were rapid and dramatic increases in photosynthetic energy conversion efficiency and phytoplankton chlorophyll concentrations (Boyd *et al.*, 2007). Beyond doubt, NPP and export production in all three regions is limited by the availability of a single micronutrient – iron.

## 2.8 Is NPP in balance with respiration?

On long time scales, a very small fraction of net primary production escapes respiration to be become buried in sediments and transferred to the lithosphere. This process has profound influence on planetary redox state. As organic matter, by definition, contains reducing equivalents, its burial requires that oxidizing equivalents accumulate elsewhere in the system. Indeed, the net burial of organic matter on geological time scales implies the oxidation of the atmosphere and ocean; i.e., the accumulation of free molecular oxygen requires burial and sequestration of organic carbon (see Chapter 7). How can we assess how much organic carbon is buried?

Recall that Rubisco strongly discriminates against the heavier isotope of carbon,  $^{13}\text{C}$ . The ultimate source of  $\text{CO}_2$  to Earth's surface is volcanism with a distinct

isotopic signal. The recycling of organic matter and carbonate rocks by weathering also represents a source of IC (inorganic carbon) to the ocean-atmosphere system. Therefore volcanism plus weathering (which should on average also carry a volcanogenic isotopic signal) represent the source of IC. The two sinks are comprised of the burial of inorganic carbonates, and organic carbon in sediments. Imagine that over some long period of time, a fraction of the total carbon input to the Earth surface is buried and preserved as organic matter. This removal of  $^{13}\text{C}$ -depleted organic carbon burial leads to an enrichment of  $^{13}\text{C}$  in the inorganic pools. The magnitude of the  $^{13}\text{C}$  carbonate enrichment depends on the proportion of the total IC input removed as organic carbon. The higher the proportion of organic carbon removal, the more enriched the  $^{13}\text{C}$  carbonates (Kump and Arthur, 1999). Removal proportions of these two fractionations are the basis of the isotopic mass balance quantified as:

$$F_{\text{in}} * \delta^{13}\text{C}_{\text{in}} = F_{\text{carb}} * \delta^{13}\text{C}_{\text{carb}} + F_{\text{org}} * \delta^{13}\text{C}_{\text{org}} \quad (2.4)$$

where  $F_{\text{in}}$  represents the total flux of IC into the ocean-atmosphere system, and  $\delta^{13}\text{C}_{\text{in}}$  is its isotopic composition. Similarly,  $F_{\text{carb}}$  and  $F_{\text{org}}$  are the burial fluxes of carbonate carbon and organic carbon, while  $\delta^{13}\text{C}_{\text{carb}}$  and  $\delta^{13}\text{C}_{\text{org}}$  are the respective isotopic compositions of these removal fluxes. At steady state, we have the equality:

$$F_{\text{in}} = F_{\text{carb}} + F_{\text{org}} \quad (2.5)$$

and furthermore, we can express the fractionation between inorganic carbon and organic carbon in the oceans as:

$$\Delta_{\text{org}} = \delta^{13}\text{C}_{\text{carb}} - \delta^{13}\text{C}_{\text{or}} \quad (2.6)$$

We also define a fraction  $f$ , which describes the burial proportion of organic carbon relative to the total flux of IC into the system

$$f = F_{\text{org}} / F_{\text{in}} \quad (2.7)$$

From these equalities, we generate the following expression for  $f$ :

$$f = (\delta^{13}\text{C}_{\text{carb}} - \delta^{13}\text{C}_{\text{in}}) / \Delta_{\text{org}} \quad (2.8)$$

At present, the average isotopic value of input DIC is ca.  $-5\%$ , the  $\delta^{13}\text{C}_{\text{carb}}$  is  $0\%$  and  $\Delta_{\text{org}}$  is ca.  $-25\%$ . Hence, to balance the input with the output, the ratio of buried organic carbon relative to total carbon is about 0.20. That is, for every 1 atom of carbon buried as organic matter, about 5 atoms are buried as carbonate (Berner, 2004). Values for  $\delta^{13}\text{C}_{\text{carb}}$  and  $\Delta_{\text{org}}$  (and probably also  $\delta^{13}\text{C}_{\text{in}}$ ) have changed through time (see Chapter 3) and hence, these serve as sedimentary archives of changes in carbon sources and sinks, thereby providing the best monitor available to reconstruct the role of the biological carbon cycle in the oxidation state of the planet's surface.

## 2.9 Conclusions and extensions

Fundamentally, the biological carbon cycle is driven by redox reactions far from thermodynamic equilibrium. On time scales of centuries and even millennia, the formation of organic matter, which is a reduction reaction, is closely coupled to respiration, an oxidation reaction, such that a quasi-steady state is achieved. The steady-state condition buffers the redox reactions on the planetary surface, and allows the highly efficient metabolic reactions of oxygen-dependent heterotrophy to proceed across vast stretches of geological time (Knoll, 2003). Indeed, animal life depends on this buffering capacity.

Humans extract fossil fuels from the lithosphere approximately 1 million times faster than they were formed. That is, each year of production of fossil fuel represents 1 million years of burial of organic matter by either plants on land (primarily yielding coal) or phytoplankton in the ocean (primarily yielding petroleum and natural gas). We burn these fossil fuels, thereby reoxidizing the organic matter back to inorganic carbon. The human combustion of fossil fuels is analogous to the action of volcanoes. However, even if humans burn the vast reserves of fossil fuels in the ground, it would have little impact on Earth's oxygen supply; there simply is so much oxygen in the atmosphere relative to the fuel available. However, the combustion of fossil fuels has rapidly altered the atmospheric inventory of  $\text{CO}_2$ , which, in turn, can alter Earth's climate and the pH of the ocean (Falkowski *et al.*, 2000). Can this problem be ameliorated by accelerating the biological carbon cycle?

The short answer is, no. Of the current emission of ca. 8.5Gt C per annum from the burning of fossil fuels, approximately 54% remains in the atmosphere; the residual 46% is split, almost evenly between terrestrial and marine ecosystems. The ability of terrestrial ecosystems to absorb the anthropogenically produced  $\text{CO}_2$  will weaken over the coming decades as the concentration of the gas begins to saturate Rubisco. Moreover, even if humans could devote large areas of land to the growth of trees, the trees would have to be harvested and preserved – they could not be allowed to enter a respiratory reaction that would oxidize the organic matter, hence the  $\text{CO}_2$  would be liberated back to the atmosphere.

In the oceans, accelerating the export of organic matter to the ocean interior would potentially help reduce the impact of anthropogenic  $\text{CO}_2$  emissions. However, the only reasonable mechanism to do this is to add a limiting nutrient to the ocean. One possible mechanism is based on the addition of iron to areas of the ocean that are iron limited. While such an approach may temporarily reduce the rate of accumulation of  $\text{CO}_2$  in the atmosphere, it is a significant geoengineering problem with potentially deleterious side-effects (Martin, 1990).

Perhaps the most interesting aspect of the biological carbon cycle is that the overwhelming majority of buried organic matter is associated with rocks, especially shales and mudstones, and is not directly accessible to oxidation on ecological time scales. The interaction between the formation of organic matter and its slow leak into the lithosphere truly distinguishes Earth from all the other planets in our solar system (Falkowski and Godfrey, 2008).

## References

- Anderson L, Sarmiento J (1994) Redfield ratios of remineralization determined by nutrient data analysis. *Global Biogeochemical Cycles* **8**, 65–80.
- Antoine D, Andre J-M, Morel A, et al. (1996) Oceanic primary production, 2: estimation at global scale from satellite (Coastal Zone Color Scanner) chlorophyll. *Global Biogeochemical Cycles* **10**(1), 57–69.
- Banse K (1992) Grazing, temporal changes of phytoplankton concentrations, and the microbial loop in the open sea. In: *Primary Productivity and Biogeochemical Cycles in the Sea* (ed Falkowski PG). Plenum, New York, pp. 409–440.
- Barber RT (1992) Geological and climatic time scales of nutrient variability. In: *Primary Productivity and Biogeochemical Cycles in the Sea* (ed Falkowski). Plenum, New York, pp. 89–106.
- Behrenfeld M, Falkowski PG (1997a) A consumer's guide to phytoplankton productivity models. *Limnology and Oceanography* **42**, 1479–1491.
- Behrenfeld MJ, Falkowski PG (1997b) Photosynthetic rates derived from satellite-based chlorophyll concentration. *Limnology and Oceanography* **42**, 1–20.
- Beja OL, Aravind EV, Koonin MT, et al. (2000) Bacterial rhodopsin: evidence for a new type of phototrophy in the sea. *Science* **289**, 1902–1906.
- Benson A, Calvin M (1950) Carbon dioxide fixation by green plants. *Annual Review of Plant Physiology* **1**, 25–42.
- Berman-Frank I, Cullen JT, Shaked Y, Sherrell RM, Falkowski PG (2001) Iron availability, cellular iron quotas, and nitrogen fixation in *Trichodesmium*. *Limnology and Oceanography* **46**, 1249–1260.
- Berner RA (2004) *The Phanerozoic Carbon Cycle: CO<sub>2</sub> and O<sub>2</sub>*. Oxford, University Press, Oxford.
- Berry J (1999) Ghosts of biospheres past. *Nature* **400**, 509–510.
- Bhattacharya D, Medlin L (1998) Algal phylogeny and the origin of land plants. *Plant Physiology* **116**, 9–15.
- Bhattacharya D, Medlin L, Wainwright PO, et al. (1992) Algae containing chlorophylls *a* + *c* are polyphyletic: molecular evolutionary analysis of the Chromophyta. *Evolution* **46**, 1801–1817.
- Blankenship RE, Madigan MT, Bauer CE, (eds) (1995) *Anoxygenic Photosynthetic Bacteria*. Kluwer Scientific, Dordrecht.
- Blankenship R, Sadekar S, Raymond J (2007) The evolutionary transition from anoxygenic to oxygenic photosynthesis. In: *Evolution of Primary Producers in the Sea* (eds Falkowski PG, Knoll AH). Academic Press, San Diego, pp. 21–35.
- Boyd PW, Jickells T, Law CS, et al. (2007) Mesoscale iron enrichment experiments 1993–2005: Synthesis and future directions. *Science* **315**, 612–617.
- Brocks JJ, Love GD, Summons RE, Knoll AH, Logan GA, Bowden SA (2005) Biomarker evidence for green and purple sulfur bacteria in an intensely stratified Paleoproterozoic sea. *Nature* **396**, 450–453.
- Broecker WS, Peng T-H (1982) *Tracers in the Sea*. EI-digio Press, Palisades, NY.
- Canfield DE (1998) A new model for Proterozoic ocean chemistry. *Nature* **396**, 450–453.
- Canfield DE, Raiswell R (1999) The evolution of the sulfur cycle. *American Journal of Science* **299**, 697–723.
- Canfield DE, Rosing MT, Bjerrum C (2006) Early anaerobic metabolisms. *Philosophical Transactions of the Royal Society B Biological Sciences* **361**, 1819–1834.
- Chisholm SW (1992) Complementary niches of *Synechococcus* and *Prochlorococcus*. In: *Primary Productivity in the Sea* (ed Falkowski PG). Plenum Press, New York, pp. 213–237.
- Codispoti L (1995) Is the ocean losing nitrate? *Nature* **376**, 724.
- de Vargas C, Probert Aubry MP, Young J (2007) Origin and evolution of coccolithophores: from coastal hunters to oceanic farmers. In: *Evolution of Primary Producers in the Sea* (eds Falkowski PG, Knoll AH). Academic Press, New York.
- DeFries R, Townshend J (1994) NDVI-derived land cover classification at global scales. *International Journal of Remote Sensing* **15**, 3567–3586.
- Delwiche C (2000) Tracing the thread of plastid diversity through the tapestry of life. *American Naturalist* **154**, S164–S177.
- Duce RA, Tindale NW (1991) Atmospheric transport of iron and its deposition in the ocean. *Limnology and Oceanography* **36**, 1715–1726.
- Falkowski PG (1997) Evolution of the nitrogen cycle and its influence on the biological sequestration of CO<sub>2</sub> in the ocean. *Nature* **387**, 272–275.
- Falkowski PG (1998) Biogeochemical controls and feedbacks on ocean primary productivity. *Science* **281**, 200–206.
- Falkowski PG (2001) Biogeochemical Cycles. *Encyclopedia of Biodiversity* **1**, 437–453.
- Falkowski PG, Godfrey L (2008) Electrons, life, and the evolution of Earth's oxygen cycle. *Philosophical Transactions of the Royal Society B Biological Sciences* **363**, 2705–2716.
- Falkowski PG, Raven JA (2007) *Aquatic Photosynthesis*. Princeton, University Press Princeton, NJ.
- Falkowski P, Barber R, Smetacek V (1998) Biogeochemical controls and feedbacks on ocean primary production. *Science* **281**, 200–206.
- Falkowski P, Scholes RJ, Boyle E, et al. (2000) The global carbon cycle: A test of our knowledge of earth as a system. *Science* **290**, 291–296.
- Falkowski P, Katz M, van Schootenbrugge B, Schofield O, Knoll AH (2003) Why is the land green and the ocean red? In: *Coccolithophores – From Molecular Processes to Global Impact* (eds Young J, Thierstein HR). Springer-Verlag, Berlin.
- Falkowski PG, Katz ME, Knoll AH, et al. (2004) The evolution of modern eukaryotic phytoplankton. *Science* **305**, 354–360.
- Fanning K (1992) Nutrient provinces in the sea: concentration ratios, reaction rate ratios, and ideal covariation. *Journal of Geophysical Research* **97C**, 5693–5712.

- Field C, Behrenfeld M, Randerson J, Falkowski P (1998) Primary production of the biosphere: integrating terrestrial and oceanic components. *Science* **281**, 237–240.
- Gruber N, Sarmiento J (1997) Global patterns of marine nitrogen fixation and denitrification. *Global Biogeochemical Cycles* **11**, 235–266.
- Han T-M, Runnegar B (1992) Megascopic eukaryotic algae from the 2.1-billion year old Neogene iron-formation. *Science* **257**, 232–235.
- Jannasch HW, Taylor CD (1984) Deep-sea microbiology. *Annual Review of Microbiology* **38**, 487–514.
- Jørgensen BB (2001) Space for hydrogen. *Nature* **412**, 286–289.
- Kaplan A, Reinhold L (1999) CO<sub>2</sub> concentrating mechanisms in photosynthetic microorganisms. *Annual Review of Plant Science and Plant Molecular Biology* **50**, 539–570.
- Kasting JF (1990) Bolide impacts and the oxidation state of carbon in the Earth's early atmosphere. *Origins Life and Evolution Biosphere* **20**, 199–231.
- Kenrick P, Crane P (1997) The origin and early evolution of plants on land. *Nature* **389**, 33–39.
- Knoll AH (1992) The early evolution of eukaryotes: a geological perspective. *Science* **256**, 622–627.
- Knoll AH (2003) *Life on a Young Planet: The First Three Billion Years of Evolution on Earth*. Princeton NJ, Princeton University Press.
- Kooistra W, Gersonde R, Medlin LK, Mann DG (2007) The origin and evolution of the diatoms: their adaptation to a planktonic existence. In: *Evolution of Primary Producers in the Sea* (eds Falkowski PG, Knoll AH). Academic Press, New York.
- Kump L, Arthur A (1999) Interpreting carbon-isotope excursions: carbonates and organic matter. *Chemical Geology* **161**, 181–198.
- Lindeman R (1942) The trophic-dynamic aspect of ecology. *Ecology* **23**, 399–418.
- Martin JH (1990) Glacial–interglacial CO<sub>2</sub> change: The iron hypothesis. *Paleoceanography* **5**, 1–13.
- Martin JH, Gordon RM, Fitzwater SE (1991) The case for iron. *Limnology and Oceanography* **36**(8), What Controls Phytoplankton Production in Nutrient-Rich Areas of the Open Sea?: 1793–1802.
- Martin W, Herrmann RG (1998) Gene transfer from organelles to the nucleus: how much, what happens, and why? *Plant Physiology* **118**, 9–17.
- Mauzerall D (1992) Light, iron, Sam Granick and the origin of life. *Photosynthesis Research* **33**(2), 163–170.
- Mooney HA, Drake BG, Luxmoore RJ, Oechel WC, Pitelka LF (1991) Predicting ecosystem responses to elevated CO<sub>2</sub> concentrations. *BioScience* **41**(2), 96–104.
- Mooney HA, Percy RW, Ehleringer J (1987) Plant physiological ecology today. *BioScience* **37**, 18–20.
- Mulkidjanian A, Junge W (1997) On the origin of photosynthesis as inferred from sequence analysis – a primordial UV-protector as common ancestor of reaction centers and antenna proteins. *Photosynthesis Research* **51**(27–42).
- Nobel PS, Forseth IN, Long SP (1993) Canopy structure and light interception. In: *Photosynthesis and Production in a Changing Environment: A Field and Laboratory Manual* (eds Hall DO, Scurlock JMO, Bolhàr-Nordenkamp HR, Leegood RC, Long SP). Chapman & Hall, London, pp. 79–90.
- Park R (1961). Carbon isotope fractionation during photosynthesis. *Geochimica et Cosmochimica Acta* **21**, 110–126.
- Postgate JR, (ed) (1971) *The Chemistry and Biochemistry of Nitrogen Fixation*. Plenum Press, New York.
- Raven JA, Falkowski PG (1999). Oceanic sinks for atmospheric CO<sub>2</sub>. *Plant Cell Environment* **22**(6), 741–755.
- Redfield A, Ketchum B, Richards F (1963) The influence of organisms on the composition of sea-water. In: *The Sea* (ed Hill M). New York, Interscience. **2**, 26–77.
- Redfield AC (1934) On the Proportions of Organic Derivatives in Sea Water and Their Relation to the Composition of Plankton. Liverpool, James Johnstone Memorial Volume.
- Redfield AC (1958) The biological control of chemical factors in the environment. *American Scientist* **46**, 205–221.
- Schlesinger WH (1997) *Biogeochemistry: An Analysis of Global Change*. Academic Press, New York.
- Shi T, Bibby TS, Jiang L, Irwin AJ, Falkowski PG (2005) Protein interactions limit the rate of evolution of photosynthetic genes in Cyanobacteria. *Molecular Biology Evolution* **22**(11), 2179–2189.
- Shi T, Falkowski PG (2008). Genome evolution in cyanobacteria: the stable core and the variable shell. *Proceedings of the National Academy of Sciences, USA* **105**, 2510–2515.
- Smith S (1981). Marine macrophytes as a global carbon sink. *Science* **211**, 838–840.
- Stevens TO, McKinley JP(1995). Lithoautotrophic microbial ecosystems in deep basalt aquifers. *Science* **270**, 450–454.
- Tyrell T (1999) The relative influences of nitrogen and phosphorus on oceanic primary production. *Nature* **400**, 525–531.
- Van Niel CB (1941) The bacterial photosyntheses and their importance for the general problem of photosynthesis. *Advances in Enzymology* **1**, 263–328.
- Warneck P (1988) *Chemistry of the Natural Atmosphere*. Academic Press, New York.
- Woese C (1998) The universal ancestor. *Proceedings of National Academy of Sciences USA* **95**: 6854–6859.
- Woese CR, Stackebrandt E, Weisburg WG, et al. (1984) The phylogeny of purple bacteria; The alpha subdivision. *Systematic and Applied Microbiology* **5**, 315–326.
- Wu JF, Sunda W, Boyle EA, Karl DM (2000) Phosphate depletion in the western North Atlantic Ocean. *Science* **289**(5480), 759–762.
- Zehr JP, Mellon M, Braun S, Litaker W, Steppe T, Paerl HW(1995) Diversity of heterotrophic nitrogen-fixation genes in a marine cyanobacterial mat. *Applied and Environmental Microbiology* **0061**, 02527–02532.
- Zehr JP, Mellon MT, Hiorns WD (1997) Phylogeny of cyanobacterial nifH genes: Evolutionary implications and potential applications to natural assemblages. *Microbiology* **143**, 1443–1450.