

# The case for Mars terraforming research

*Can we understand enough about climate and ecosystems to build them elsewhere?*

Erika Alden DeBenedictis<sup>1</sup>, Edwin S. Kite<sup>2</sup>, Robin D. Wordsworth<sup>3</sup>, Nina L. Lanza<sup>4</sup>, Charles S. Cockell<sup>5</sup>, Pamela A. Silver<sup>6</sup>, Ramses M. Ramirez<sup>7</sup>, John Cumbers<sup>8</sup>, Hooman Mohseni<sup>9</sup>, Christopher E. Mason<sup>10</sup>, Woodward W. Fischer<sup>11</sup>, & Christopher P. McKay<sup>12</sup>.

1. Pioneer Research Labs (erika@pioneer-labs.org), 2. University of Chicago, 3. Harvard University, 4. Los Alamos National Laboratory, 5. University of Edinburgh, 6. Harvard - Wyss, 7. University of Central Florida, 8. SynBioBeta, 9. Northwestern University, 10. Weill Cornell Medicine, 11. Caltech, 12. NASA Ames Research Center.


Exploration of Mars is motivated by scientific, societal, and engineering goals, and the centuries-old dream that people could one day live there. Missions to date reveal a hostile surface environment, featuring a deadly triple cocktail of extreme cold, high surface UV fluxes, and lack of sustained liquid water [1]. Despite this harshness, advocacy for large-scale Mars settlement has continued for decades, but is making Mars' surface hospitable to life possible and feasible?

Proposed motivations for terraforming are diverse. Carl Sagan once wrote, "If we do not destroy ourselves, we will one day venture to the stars." Since Sagan's time we have confirmed that Mars was once habitable but suffered a global climate catastrophe, so some view Mars as an environmental restoration challenge. Some argue that a hospitable Mars is essential to achieve self-sufficiency, surpassing the limitations of isolated outposts. Others are motivated by the scientific desire to learn about the universe, as the realization of humanity's dreams to explore the universe is assisted by expanded human presence.

Contemplating building a climate and ecosystem beyond Earth highlights gaps in our fundamental knowledge of these systems. Ongoing developments in three key areas have returned terraforming to the martian research agenda. First, our understanding of climate modeling and climate engineering, including for Earth, has advanced. It is timely to investigate how these techniques might be applied on other planets. Second, progress in synthetic biology has improved our knowledge of extremophilic organisms and our ability to engineer their properties, opening new possibilities for tailoring life to thrive in the extremes of Mars. Third are numerous developments in space science. The emergence of Mars transport vehicles like Starship will broaden the scope of possible space missions by increasing the mass we can launch from Earth by >100× per Mars landing [2]. Our understanding of the basic science of Mars has advanced, leading to consensus that a warmed Mars would retain volatiles for many millions of years [3]. Finally, new options for warming Mars have emerged, using ultralight materials [4], solar sails [5], or nanoparticles [6]. Thus, a fresh look at the research agenda for greening Mars [7-8] is timely. Restoring a habitable planet is harder than sustaining one, and thus Mars presents the ultimate sustainability challenge.

We consider three thresholds in making Mars suitable for life (Fig 1). Each threshold motivates complementary research programs that should start now. The first

phase in greening Mars would involve abiotic environmental engineering to heat the planet, locally and/or globally. A future, warmer Mars would pass the threshold of creating conditions over large regional scales suitable for extreme life. In the second phase, some extreme species will be able to grow within the life-compatible area. As the first step, an autotrophic, (likely) anaerobic primary producer is needed, followed by ecological succession to diversify and stabilize the ecosystem [9]. A green planet with a flourishing biosphere including algae and plants would constitute a second threshold. The third phase aims at developing a biosphere with complex plant life and perhaps trees (but not necessarily animal life), with increased  $O_2$  content and atmospheric pressure. We now consider each phase in turn.



### Short term.

Mars' thin atmosphere (~6 mbar  $CO_2$ ) results in a global average temperature of  $-70^\circ C$  (despite local highs exceeding  $20^\circ C$ ), and precludes stable surface liquid water across much of the planet, limiting biological potential. Increasing liquid water availability would enable scalable biological techniques for agriculture and ecological succession, improving Mars's habitability for humans.

Mars has less  $H_2O$  than Earth, but still has at least enough ice to form a  $>300$  m deep ocean over  $10^7$   $km^2$  of the planet [11]. However, known  $H_2O$  reserves are in the


coldest third of Mars's surface [12], with minimal confirmed water in warmer regions ( $2\pm 1$  wt% water-equivalent hydrogen in minerals, largely releasable at  $<350$  °C [13]). Thus, for Mars to be livable, temperature must rise to the point where melting starts at relatively cold locations. These H<sub>2</sub>O-ice-bearing locations are already prioritized for human missions as H<sub>2</sub>O can be processed for propellant, and their temperature likely needs to increase by at least 30°C to start to melt the ice. An added challenge for melting is that evaporitic cooling hinders ice melting on Mars, because of the large amount of energy required to transform ice into vapor [14].

Warming Mars could be achieved by increasing insolation (currently 130 W/m<sup>2</sup>) using solar sails as mirrors to collect and reflect additional light onto the planet [5], or enhancing the greenhouse effect. Local enhancement methods include tiling the surface with silica aerogels (solid-state greenhouse effect) or nanocellulose [4,14], while regional/global approaches include engineered aerosols [5]. These techniques appear much more mass-effective than earlier proposals based on anthropogenic atmospheric warming on Earth (using fluorocarbon gases [15]). Mars' low atmospheric thermal inertia ensures faster (global or local) warming responses to radiative balance changes compared to Earth.

Further research is crucial to model warming's effects on Mars's climate. Recent advances in Earth's climate models can be applied to Mars; similarly, the challenge of expanding these models to a new planet could improve predictions for both. Many effects must be carefully modeled to generate realistic predictions. Warming will perturb water, CO<sub>2</sub>, and dust cycles [10, 16], reactivating feedbacks from wetter periods in Mars history and that could help or hinder near-future warming efforts. Atmospheric thickness will at least double as buried CO<sub>2</sub> ice is released [17]. While H<sub>2</sub>O-vapor feedback is positive and cloud feedback likely positive (but of uncertain magnitude) [18], dust cycle intensification may warm the planet overall but lower peak temperatures [19]. Research is needed to track ground ice redistribution (toward the equator, or onto high ground?) as the water cycle intensifies [20-21] and to simulate dust cycle changes under warmer conditions [22]. Model intercomparisons are also needed [23].

In addition, we must raise the Technology Readiness Level (TRL) of proposed warming approaches. The path to deployment is long, and requires forethought. For example, engineered aerosols require laboratory validation of key microphysical parameters, and wind-tunnel studies. A small-scale field test within Mars' atmosphere would be needed to validate models, in turn requiring greenhouse-agent plume dispersal calculations and also monitoring instruments (which can also be used to enhance Mars weather/climate science). Any warming method (local or global) must be controllable, reversible within years, and biocompatible, factors that will shape deployment strategies [24].

Together, advances in Earth's launch capacity, combined with proposed new warming techniques, could potentially raise Mars' temperature by 30°C well within the century, permitting liquid H<sub>2</sub>O for the first living organisms to grow on the surface.



### Mid-term.

A warmer Mars could support extreme life, initiating ecological succession towards a diverse ecosystem that begins producing an oxygen-rich atmosphere [25-26].

Engineering pioneer species capable of growth (assuming the presence of liquid water) despite Mars' unique mixture of five primary stressors - low pressure, oxychlorine species, low temperature, radiation (including ionizing and ultraviolet light), and low water activity (caused, for example, by high ionic concentrations in many locations) (Figure 3) - may be achievable. 1) The pioneer species would need to be anaerobic and tolerate low atmospheric pressures. Microbes can grow at Mars-like pressures of 7 mbar [27]. 2) Oxychlorine salts like perchlorate and chlorate are widespread on Mars. Fortunately, some  $\text{O}_2$ -generating microbes can use perchlorate reduction for metabolism [28]. 3) The pioneer species will need to grow at cold temperatures, with large day-night temperature swings. Organisms exist that can grow at  $-12^\circ \text{ C}$  [29] and tolerate daily freeze-thaw [30]. 4) Mars radiation (while a problem for humans) is not a problem for microbes. UV-C levels on Mars are high ( $\sim 3 \text{ W/m}^2$  of UV-C), but can be screened sufficiently while still allowing


124 phototrophy [31]. 5) As Mars is heated, the first liquid water will be salty brines including  
125 mixtures of chloride and sulfate anions, potentially requiring halophilic microbes, which  
126 are plentiful on Earth.

127 Other Martian conditions are potentially suitable for microbial growth. These  
128 include Mars gravity, which has been shown to be consistent with microbial growth [32].  
129 Although the atmosphere lacks significant N<sub>2</sub>, the soil contains fixed nitrogen phases,  
130 principally nitrate/nitrite (110–300 ppmw) [33]. The pioneer organism's home - whether a  
131 surface pond, or ground water - will gain nutrients from Mars soil [34-35]. The soil at  
132 lander sites can (by Earth standards) have high ionic concentrations but be quite nutrient-  
133 rich, and ½ kg/m<sup>3</sup> of organic carbon is reported from some Mars sediments [36]. Soil pH  
134 is 7.7±0.3 at 68°N [37]. Phosphate release rates during water–rock interactions on Mars  
135 are thought to be much higher than on Earth [38]. Mars-like rocks (volcanic basalt) support  
136 diverse pioneering microbial communities in Mars-like regions of Earth [39].

137 Mars-adapted organisms may be developed through genetic engineering, directed  
138 evolution [40], and Mars-chamber experiments with extremophiles. This research has  
139 much overlap with existing science priorities. For example, soil sample return would refine  
140 the target habitat by allowing quantification of biocritical trace elements and possible  
141 toxins [41]. This research overlaps with Earth green biomanufacturing priorities [42].

142 Candidate patchy water deposits at Mars' equator (e.g., [43]) could be watering  
143 holes well-suited to attract photosynthetic organisms and (inside protective membranes)  
144 people. At this stage, we envisage microbes supporting the food and oxygen needs for  
145 10<sup>4</sup> people per site [44], but with automation assisting with gardening/farming (the outside  
146 atmosphere will not be breathable). O<sub>2</sub> from perchlorate reduction and photosynthesis  
147 could be initially confined within production environments. Once O<sub>2</sub> is sufficient to support  
148 respiration in plants (and/or humans) within local membrane-bound environments, excess  
149 can be gradually released to the global atmosphere.

150 By the end of the second phase of terraforming, extremophiles would be at work  
151 to transform planetary chemistry. A warmed Mars with an oxygen- and food-producing  
152 ecosystem could allow many more people to live on Mars, and would lead the way for a  
153 more complex ecosystem and thicker atmosphere.



**Terraforming and the search for extant Martian life** The idea of terraforming Mars emphasizes the motivation to support people living beyond Earth. There is some reluctance among planetary scientists to contemplate adding more Earth-derived life to Mars (beyond what is already added by landed spacecraft [45]), because Earth microbes might be confused for (or supplant) Mars-derived life, if it exists. This is mitigated by the recognition that landing humans on Mars will introduce orders of magnitude more Earth microbes to the Mars environment [45]. Therefore, with humans living on Mars in the near future, a concerted effort is needed to determine if Mars has life. This could be greatly aided by human outposts, and should include soil sample return, and sounding for deep aquifers. If life is detected, then its scientific importance could warrant robust protections for its habitat.

### Long term.

Once Mars possesses a planetary ecosystem, it would continue to develop over the long-term. It has been argued that the long-term goal for astrobiology should be to enhance the richness and diversity of life in the Universe [47]. What would a target atmosphere for Mars look like? Key *in situ* sources of volatiles include  $\text{H}_2\text{O}$  and  $\text{CO}_2$ .  $\text{H}_2\text{O}$  can be used to create  $\text{O}_2$ .  $\text{CO}_2$  will be released naturally as the planet warms, and can also be extracted from carbonates - but high levels of  $\text{CO}_2$  are toxic to humans [48]. Notably, Mars lacks an obvious source of *in situ* inert gas that might take the role of  $\text{N}_2$  in Earth's atmosphere. A 0.1 bar  $\text{O}_2$  atmosphere could satisfy human habitability requirements, including breathability and reducing Mars surface radiation to Earth-airliner levels.  $\text{O}_2$  levels cannot rise too high however due to the growing risks of uncontrolled combustion. Together, a target atmosphere containing 100 mbar  $\text{O}_2$  satisfies all these requirements: feasible to

168 achieve entirely with *in situ* atoms and suitable for human habitation. This atmosphere  
169 could be generated within 1-2 years in 100-meter tall domes using photosynthesis or  
170 water electrolysis [49].

171 In addition to domed habitats suitable for humans without pressure suits, more  
172 species could inhabit the surface, albeit at lower pressure. As O<sub>2</sub> builds up, more species  
173 could live on the surface, and the fraction of (potentially tented/domed) Mars surface area  
174 where humans could breathe would increase. An intriguing possibility is self-extending  
175 (similar to coral reefs) O<sub>2</sub>-impermeable membranes produced by life [4, 50]. Organisms  
176 or their biofilms might also modulate planetary energy balance through albedo effects and  
177 solid-state greenhouse warming [4, 51].

178 While rapid greening of Mars may be possible, establishing a global 0.1 bar O<sub>2</sub>  
179 atmosphere through photosynthesis alone would take millennia. Oxygenation via  
180 photosynthesis would involve complex biogeochemical cycles [52,53], including O<sub>3</sub> shield  
181 formation and organic matter sequestration. Once a favorable climate is created on Mars,  
182 it opens new questions, such as how to stabilize and maintain that climate. Historically,  
183 Mars has experienced relatively wetter and drier periods, at least regionally, with its  
184 extensive high-elevation surfaces that trap water as high albedo ice caps being a major  
185 factor [20]. Climate stabilization would require understanding water movement between  
186 oceans, high-elevation ice, and subsurface aquifers (and its impact on albedo and thus  
187 temperature), as well as potential water loss to the deep subsurface through taliks (layers  
188 of unfrozen water in permafrost regions).

189 Current research needs include improved climate models and more spacecraft  
190 data to assess long-term climate possibilities. Future missions [46] to determine what lies  
191 beneath the ground ice that is known to exist over ~½ of Mars's surface - liquid water, or  
192 empty pore space, or more water ice - will be important for setting how large Martian lakes  
193 and seas can be. After the use of light to split water is well-established, oxygen build-up  
194 will require electron acceptors. Quantifying reservoirs of electron acceptors - including  
195 sulfates/Fe-oxides, and CO<sub>2</sub> from various sources ( $\leq 100$  mbar [54-55]) or carbonate - is  
196 essential, as if electron acceptors are in short supply then it will not be possible for  
197 humans to breathe unaided on the surface without expensive importation of volatiles from  
198 beyond Mars. It has long been recognized that currently unknown surface and subsurface  
199 reservoirs of CO<sub>2</sub>, H<sub>2</sub>O and nitrate are key to enabling a global biosphere on Mars [8,54].  
200 Discoveries are ongoing: in 2023, the *Curiosity* rover discovered abundant carbonate not  
201 seen from orbit.

202 If this stage succeeds, the planet would have a stable, favorable climate, and  
203 support a diverse planetary biosphere. The outcome would be something new and  
204 different - not a replacement for Earth, but an addition.

## 205 206 **The futures of Mars.**

207  
208 New ideas for Mars' future have emerged that are sustainable, resilient, and ecologically  
209 minded [56]. Technologies developed for Mars habitation, such as desiccation-resistant  
210 crops, efficiently remediating soil, and improved ecosystem modeling [9], will likely benefit  
211 Earth. Rather than distracting from Earth's problems, Mars terraforming research could  
212 provide insights for maintaining oasis Earth, expanding rather than diminishing our  
213 environmental responsibility. Mars terraforming research offers a vital testbed for

planetary science, potentially validating theories or exposing knowledge gaps. Continued research promises significant scientific progress, regardless of whether full-scale terraforming occurs.

While the possibilities are exciting, anything as big as modification of a planetary climate has major consequences, and would require careful thought once we reached the point where it was feasible. But until we do more research, we don't even know what's physically or biologically possible. Therefore, further Mars exploration is crucial. Priorities include quantifying H<sub>2</sub>O, N<sub>2</sub>, and CO<sub>2</sub> reserves (e.g., searching for deep aquifers [46]), soil sample return, test missions for proof-of-concept of warming methods, and climate feedback studies. These align with existing mission priorities [57,41]: for example, ice deposit investigations [58] constrain the extent of resources whose abundance could fuel (or restrict) terraforming, and geologically-recent warm climates [18,59] are a natural analog for a near-future warmed Mars. Current Mars Exploration Program Analysis Group goals already support human habitation [57]. No abrupt change of course is needed: indeed, support for people living on Mars provides fresh rationale for many existing mission priorities, alongside precursor/test missions.

Beyond our Solar System, rocky planets are common [60], but worlds that are ready for life will be rare. If people can learn how to terraform a world such as Mars, this may be the first step to destinations beyond. More speculatively, the technologies eventually determined to best enable terraforming will refine our ability to search for technosignatures [61].

When people start to live on Mars, "they will inevitably introduce orders of magnitude more terrestrial microorganisms to Mars than robotic missions have done or will do" [45]. The open question is whether we engage with Mars in an informed way, or an uninformed way. As large corporations and governments contemplate Mars terraforming, we suggest that science must have an important role to play. This is only possible if research accelerates appropriately to keep pace with Mars-access capabilities.

**Acknowledgements.** We thank The Astera Institute for support of a workshop. We thank all of the workshop attendees for creating a vibrant brainstorming environment.

## References.

- [1] Martín-Torres, F. J., Zorzano, M. P., Valentín-Serrano, P., Harri, A. M., Genzer, M., Kemppinen, O., ... & Vaniman, D. (2015). Transient liquid water and water activity at Gale crater on Mars. *Nature Geoscience*, 8(5), 357-361.
- [2] Heldmann, J.L., Marinova, M.M., Lim, D.S., Wilson, D., Carrato, P., Kennedy, K., Esbeck, A., Colaprete, T.A., Elphic, R.C., Captain, J. and Zacny, K., 2022. Mission architecture using the SpaceX Starship vehicle to enable a sustained human presence on Mars. *New Space*, 10(3), pp.259-273.
- [3] Jakosky, B.M. "Mars' atmosphere, volatiles, and climate as the sun heats up over the next 6 billion years." *Icarus* 410 (2024): 115888.
- [4] Wordsworth, R., Kerber, L., & Cockell, C. (2019). Enabling Martian habitability with silica aerogel via the solid-state greenhouse effect. *Nature Astronomy*, 3(10), 898-903.
- [5] Handmer, C. (2024), How to terraform Mars for \$50bn with solar sails, *Tenth International Conference on Mars*, Pasadena, CA, abstract #3025.
- [6] Ansari, S., Kite, E.S., Ramirez, R., Steele, L., & Mohseni, H. (2024), Feasibility of keeping Mars warm with nanoparticles, *Science Advances*, 10(32), p.eadn4650.



- [7] Avernier, M. M., & MacElroy, R. D. (1976). *On the habitability of Mars: An approach to planetary ecosynthesis* (NASA-SP-414).
- [8] McKay, C.P., Toon, O.B. and Kasting, J.F., 1991. Making Mars habitable. *Nature*, 352(6335), pp.489-496.
- [9] Goyal, A., Flamholz, A.I., Petroff, A.P. and Murugan, A., 2023. Closed ecosystems extract energy through self-organized nutrient cycles. *Proc. Natl. Acad. Sci.*, 120(52), p.e2309387120.
- [10] Haberle, R. M., Clancy, R. T., Forget, F., Smith, M. D., & Zurek, R. W. (Eds.). (2017). *The atmosphere and climate of Mars*. Cambridge University Press.
- [11] Carr, M. H., & Head, J. W. (2015). Martian surface/near-surface water inventory: Sources, sinks, and changes with time. *Geophys. Res. Lett.*, 42(3), 726-732.
- [12] Morgan, G. A., Putzig, N. E., Perry, M. R., Sizemore, H. G., Bramson, A. M., Petersen, E. I., ... & Campbell, B. A. (2021). Availability of subsurface water-ice resources in the northern mid-latitudes of Mars. *Nature Astronomy*, 5(3), 230-236.
- [13] Leshin, L. A., Mahaffy, P. R., Webster, C. R., Cabane, M., Coll, P., Conrad, P. G., ... & Fabre, C. (2013). Volatile, isotope, and organic analysis of martian fines with the Mars Curiosity rover. *Science*, 341(6153), 1238937.
- [14] Schorghofer, N. (2020). Mars: Quantitative evaluation of crocus melting behind boulders. *The Astrophysical Journal*, 890(1), 49.
- [14] Smalyukh, I.I., 2021. Thermal management by engineering the alignment of nanocellulose. *Advanced Materials*, 33(28), p.2001228.
- [15] Marinova, M. M., McKay, C. P., & Hashimoto, H. (2005). Radiative-convective model of warming Mars with artificial greenhouse gases. *J. Geophys. Res.: Planets*, 110(E3).
- [16] Bertrand, T. R. J. Wilson, M. A. Kahre, R. Urata, and A. Kling. Simulation of the 2018 global dust storm on Mars using the NASA Ames Mars GCM: a multitracer approach. *J. Geophys. Res. - Planets* 125, no. 7 (2020): e2019JE006122.
- [17] Bierson, C. J., Phillips, R. J., Smith, I. B., Wood, S. E., Putzig, N. E., Nunes, D., & Byrne, S. (2016). Stratigraphy and evolution of the buried CO<sub>2</sub> deposit in the Martian south polar cap. *Geophys. Res. Lett.*, 43(9), 4172-4179.
- [18] Madeleine, J.B., Head, J.W., Forget, F., Navarro, T., Millour, E., Spiga, A., Colaïtis, A., Määttänen, A., Montmessin, F. and Dickson, J.L., 2014. Recent ice ages on Mars: the role of radiatively active clouds and cloud microphysics. *Geophys. Res. Lett.*, 41(14), pp.4873-4879.
- [19] Streeter, P. M., Lewis, S. R., Patel, M. R., Holmes, J. A., & Kass, D. M. (2020). Surface warming during the 2018/Mars Year 34 global dust storm. *Geophys. Res. Lett.*, 47(9), e2019GL083936.
- [20] Wordsworth, R. D., Kerber, L., Pierrehumbert, R. T., Forget, F., & Head, J. W. (2015). Comparison of “warm and wet” and “cold and icy” scenarios for early Mars in a 3-D climate model. *J. Geophys. Res.: Planets*, 120(6), 1201-1219.
- [21] Ramirez, R. M., & Craddock, R. A. (2018). The geological and climatological case for a warmer and wetter early Mars. *Nature Geoscience*, 11(4), 230-237.
- [22] Green, J. L., Airapetian, V., Bamford, R., Call, N., Dong, C., Hollingsworth, J., ... & Wheeler, R. (2021). Interdisciplinary Research in Terraforming Mars: State of the Profession and Programmatics. National Academy of Sciences Decadal Survey White Paper white paper; Bulletin of the American Astronomical Society, Vol. 53, Issue 4, e-id. 488
- [23] Fauchez, T. J., Turbet, M., Wolf, E. T., Boutle, I., Way, M. J., Del Genio, A. D., ... & Domagal Goldman, S. D. (2020). TRAPPIST-1 Habitable Atmosphere Intercomparison (THAI): motivations and protocol version 1.0. *Geoscientific Model Development*, 13(2), 707-716.
- [24] MacMartin, D. G., & Kravitz, B. (2019). The engineering of climate engineering. *Annual Review of Control, Robotics, and Autonomous Systems*, 2, 445-467.
- [25] Graham, J. M. (2004). The biological terraforming of Mars: planetary ecosynthesis as ecological succession on a global scale. *Astrobiology*, 4(2), 168-195.

- [26] Cyclic, L. M., Hausrath, E. M., Ming, D. W., Adcock, C. T., Raymond, J., Remias, D., & Ruemmele, W. P. (2021). Investigating the growth of algae under low atmospheric pressures for potential food and oxygen production on Mars. *Frontiers in Microbiology*, 12, 733244.
- [27] Schuerger, A. C., & Nicholson, W. L. (2016). Twenty species of hypobarophilic bacteria recovered from diverse soils exhibit growth under simulated martian conditions at 0.7 kPa. *Astrobiology*, 16(12), 964-976.
- [28] Davila, A. F., Willson, D., Coates, J. D., & McKay, C. P. (2013). Perchlorate on Mars: a chemical hazard and a resource for humans. *International Journal of Astrobiology*, 12(4), 321-325.
- [29] Breezee, J., Cady, N. and Staley, J.T., 2004. Subfreezing growth of the sea ice bacterium "Psychromonas ingrahamii". *Microbial Ecology*, 47, pp.300-304.
- [30] Walker, V.K., Palmer, G.R. and Voordouw, G., 2006. Freeze-thaw tolerance and clues to the winter survival of a soil community. *Applied and Environmental Microbiology*, 72(3), pp.1784-1792.
- [31] Cockell, C. S., & Raven, J. A. (2004). Zones of photosynthetic potential on Mars and the early Earth. *Icarus*, 169(2), 300-310.
- [32] Lima, I. G. P., McCutcheon, G., Kent, R., Pless, E. E., Ricco, A. J., White, B., ... & Rothschild, L. (2021). PowerCell aboard the EuCROPIS satellite-Results from the first synthetic biology experiment in space. In *The American Society for Gravitational and Space Research (ASGSR) 2021 meeting*.
- [33] Stern, J.C., Sutter, B., Freissinet, C., Navarro-González, R., McKay, C.P., Archer Jr, P.D., Buch, A., Brunner, A.E., Coll, P., Eigenbrode, J.L. and Fairen, A.G., 2015. Evidence for indigenous nitrogen in sedimentary and aeolian deposits from the Curiosity rover investigations at Gale crater, Mars. *Proc. Natl. Acad. Sci.*, 112(14), pp.4245-4250.
- [34] O'Connell-Cooper, C. D., Spray, J. G., Thompson, L. M., Gellert, R., Berger, J. A., Boyd, N. I., ... & VanBommel, S. J. (2017). APXS-derived chemistry of the Bagnold dune sands: Comparisons with Gale Crater soils and the global Martian average. *J. Geophys. Res.: Planets*, 122(12), 2623-2643.
- [35] Sutter, B., McAdam, A. C., & Mahaffy, P. R. (2019). Volatile detections in Gale crater sediment and sedimentary rock: results from the Mars Science Laboratory's sample analysis at Mars instrument. In *Volatiles in the Martian Crust* (pp. 369-392). Elsevier.
- [36] Stern, J. C., Malespin, C. A., Eigenbrode, J. L., Webster, C. R., Flesch, G., Franz, H. B., ... & Mahaffy, P. R. (2022). Organic carbon concentrations in 3.5-billion-year-old lacustrine mudstones of Mars. *Proc. Natl. Acad. Sci.*, 119(27), e2201139119.
- [37] Kounaves, S. P., & Oberlin, E. A. (2019). Volatiles measured by the phoenix lander at the northern plains of Mars. In *Volatiles in the Martian Crust* (pp. 265-283). Elsevier.
- [38] Adcock, C. T., Hausrath, E. M., & Forster, P. M. (2013). Readily available phosphate from minerals in early aqueous environments on Mars. *Nature Geoscience*, 6(10), 824-827.
- [39] Kelly, L.C., Cockell, C.S., Thorsteinsson, T., Marteinson, V. and Stevenson, J. (2014) Pioneer microbial communities of the Fimmvörðuháls lava flow, Eyjafjallajökull, Iceland. *Microbial Ecology* 68, 504-518.
- [40] Menezes, A. A., Cumbers, J., Hogan, J. A., & Arkin, A. P. (2015). Towards synthetic biological approaches to resource utilization on space missions. *Journal of the Royal Society Interface*, 12(102), 20140715.
- [41] National Research Council, Space Studies Board, Space Engineering Board, & Committee on Precursor Measurements Necessary to Support Human Operations on the Surface of Mars. (2002). *Safe on Mars: Precursor measurements necessary to support human operations on the Martian surface*. National Academies Press.
- [42] DeBenedictis, E., Caro-Astorga, J., Stork, D., Nattermann, U., Piskiewicz, S., Vimercati, L., Schwendner, P., Meyerowitz, J.T., Hoher, A. and Cockell, C., (2024), Polyextremophile

engineering: a review of organisms that push the limits of life. *Frontiers in Microbiology*, 15, p.1341701.

[43] Shean, D. E. (2010). Candidate ice-rich material within equatorial craters on Mars. *Geophys. Res. Lett.*, 37(24).

[44] Nangle, S. N., Wolfson, M. Y., Hartsough, L., Ma, N. J., Mason, C. E., Merighi, M., ... & Ziesack, M. (2020). The case for biotech on Mars. *Nature Biotechnology*, 38(4), 401-407.

[45] Stern, A., et al., (2019), Final Report of the NASA Planetary Protection Independent Review Board, downloaded from [https://www.nasa.gov/wp-content/uploads/2015/01/planetary\\_protection\\_board\\_report\\_20191018.pdf?emrc=9b0ac1](https://www.nasa.gov/wp-content/uploads/2015/01/planetary_protection_board_report_20191018.pdf?emrc=9b0ac1)

[46] McGarey, P., Carpenter, K., Nunes, D., Barba, N., Manthena, R., Krieger, S., ... & Stamenkovic, V. (2022, March). Development of a Loop Antenna Deployment System for TEM-Based Subsurface Mars Water Detection. In *2022 IEEE Aerospace Conference (AERO)* (pp. 1-11). IEEE.

[47] McKay C., (2013), in Impey et al., eds., *Encountering Life in the Universe*, University of Arizona Press, p. 158-166.

[48] Schwieterman, E. W., Reinhard, C. T., Olson, S. L., Harman, C. E. & Lyons, T. W. A limited habitable zone for complex life. *Astrophys. J.* 878, 19 (2019).

[49] West, W., Atwater, H. A., & Kubiak, C. (2018). Addressing the Mars ISRU Challenge: Production of Oxygen and Fuel from CO<sub>2</sub> Using Sunlight. *Keck Institute for Space Studies*.

[50] Wordsworth, R., & Cockell, C. Defining the minimum physical requirements for life in space, in review.

[51] Lenton, T. M., & Lovelock, J. E. (2001). Daisyworld revisited: quantifying biological effects on planetary self-regulation. *Tellus B*, 53(3), 288-305.

[52] Fischer, W. W., Hemp, J., & Johnson, J. E. (2016). Evolution of oxygenic photosynthesis. *Annual Review of Earth and Planetary Sciences*, 44, 647-683.

[53] Knoll, A. H., Canfield, D. E., & Konhauser, K. (Eds.). (2012). *Fundamentals of geobiology* (pp. 297-314). Chichester: Wiley-Blackwell.

[54] Jakosky, B. M., & Edwards, C. S. (2018). Inventory of CO<sub>2</sub> available for terraforming Mars. *Nature astronomy*, 2(8), 634-639.

[55] Buhler, P. B., & Piqueux, S. (2021). Obliquity-Driven CO<sub>2</sub> Exchange Between Mars' Atmosphere, Regolith, and Polar Cap. *J. Geophys. Res.: Planets*, 126(5), e2020JE006759.

[56] Turner, F., Cultivating a planet - and ourselves, *Harpers*, August 1989, 33-40

[57] MEPAG (2020), Mars Scientific Goals, Objectives, Investigations, and Priorities: 2020. D. Banfield, ed., 89 p. white paper posted March, 2020 by the Mars Exploration Program Analysis Group (MEPAG) at <https://mepag.jpl.nasa.gov/reports.cfm>

[58] Smith, I. B., Hayne, P. O., Byrne, S., Becerra, P., Kahre, M., Calvin, W., ... & Siegler, M. (2020). The Holy Grail: A road map for unlocking the climate record stored within Mars' polar layered deposits. *Planetary and space science*, 184, 104841.

[59] Dickson, J.L., Palumbo, A.M., Head, J.W., Kerber, L., Fassett, C.I. and Kreslavsky, M.A., 2023. Gullies on Mars could have formed by melting of water ice during periods of high obliquity. *Science*, 380(6652), pp.1363-1367.

[60] Bryson, S., Kunimoto, M., Kopparapu, R.K., Coughlin, J.L., Borucki, W.J., Koch, D., Aguirre, V.S., Allen, C., Barentsen, G., Batalha, N.M. and Berger, T., 2020. The occurrence of rocky habitable-zone planets around solar-like stars from Kepler data. *The Astronomical Journal*, 161(1), p.36.

[61] Wright, J. T., Haqq-Misra, J., Frank, A., Kopparapu, R., Lingam, M., & Sheikh, S. Z. (2022). The case for technosignatures: Why they may be abundant, long-lived, highly detectable, and unambiguous. *The Astrophysical Journal Letters*, 927(2), L30.

[62] Paulino-Lima, I.G., Fujishima, K., Navarrete, J.U., Galante, D., Rodrigues, F., Azua-Bustos, A. and Rothschild, L.J., 2016. Extremely high UV-C radiation resistant microorganisms from

412 desert environments with different manganese concentrations. *Journal of Photochemistry and*  
413 *Photobiology B: Biology*, 163, pp.327-336.  
414 [63] Heinz, J., Krahn, T. and Schulze-Makuch, D., 2020. A new record for microbial perchlorate  
415 tolerance: fungal growth in NaClO<sub>4</sub> brines and its implications for putative life on Mars. *Life*, 10(5),  
416 p.53.